


Redox & Electrochemistry Practice Items

1. An oxidizing agent
- A. receives electrons in a redox reaction.
 - B. supplies electrons in a redox reaction.
 - C. tends to contain atoms with low oxidation numbers.
 - D. reacts spontaneously with O_2 .

2. What is the oxidation number of chlorine in dichlorine heptoxide?




- A. -4
 - B. 0
 - C. +4
 - D. +7
3. Which of the following statements is true regarding the reaction of copper(II) sulfate with iron?




- I. Iron is reduced
 - II. Copper is reduced
 - III. Sulfate ion serves as an oxidizing agent
- A. I only
 - B. II only
 - C. I and III
 - D. II and III

4. Which of the following is true regarding molecular oxygen and molecular hydrogen?
- A. O_2 is usually reduced in reaction (not with fluorine), while H_2 is usually oxidized (though not by metals).
 - B. The oxidation number of oxygen in O_2 is -2, while the oxidation number of hydrogen in H_2 is +1.
 - C. O_2 is a strong reducing agent, while H_2 is usually an oxidizing agent.
 - D. Hydrogen gas production from a reaction vessel is often the result of the reduction of a metal.
5. Which of the following metals will react vigorously with liquid water to give H_2 ?
- A. Ag
 - B. Au
 - C. Hg
 - D. L
6. Potassium permanganate, $KMnO_4$, is a strong oxidizing agent. In reactions with potassium permanganate, which atom is reduced?
- A. K
 - B. Mn
 - C. O
 - D. The substrate is reduced.
7. Many pure metals can be produced by reaction of their chlorides with sodium because
- A. sodium forms an ionic compound with chlorine.
 - B. sodium ion possesses a very high negative standard reduction potential.
 - C. sodium is a very electrophilic element.
 - D. sodium chloride has a large lattice energy.

8. What is the coefficient on chlorine molecule when the following oxidation reduction reaction (in acidic solution) is balanced?



- A. 1
B. 3
C. 6
D. 7
9. In the reaction below, which species is oxidized and which is reduced?



- A. H is oxidized and N is reduced.
B. N is oxidized and O is reduced.
C. N is both oxidized and reduced.
D. N is oxidized and H is reduced.
10. A large, positive standard reduction potential for a substance indicates that the substance is
- A. a strong reducing agent.
B. a strong oxidizing agent.
C. easily oxidized.
D. a cation.

11. In an electrolytic cell

- A. oxidation occurs at the positive anode.
B. reduction occurs at the positive anode.
C. oxidation occurs at the negative anode.
D. reduction occurs at the negative anode.

12. In a galvanic cell


- A. oxidation occurs at the positive anode.
B. reduction occurs at the positive anode.
C. oxidation occurs at the negative anode.
D. reduction occurs at the negative anode.

13. Which of the following results from the electrolysis of a solution of 5M sodium chloride?

- I. production of sodium at the anode and chlorine gas at the cathode
II. production of chlorine gas at the anode and hydrogen gas at the cathode
III. a basic solution around the cathode

- A. I only
B. II only
C. I and III
D. II and III

14. The anode and cathode reactions are as follows:



The standard reduction potential of Zn^{2+} is -0.762 V , and that of Ag^+ is $+0.800\text{ V}$. What is the approximate emf of a silver oxide battery?

- A. 0.04 V
B. 0.8 V
C. 1.6 V
D. 2.4 V

15. A battery runs dead when the redox reaction
- has moved charge through the salt bridge equaling capacitance times voltage.
 - has consumed all available reagents.
 - has led to creation of an equal and opposite potential in the salt bridge.
 - reaches the equilibrium state.

16. Placing metallic zinc into a solution of 1M CuCl_2 will result in a layer of copper being deposited on the zinc. Placing an aluminum rod in a 1M solution of ZnCl_2 will result in metallic zinc being deposited on the aluminum. Which of the following can be concluded from these observations?

- Zinc is a stronger reducing agent than copper.
- The standard reduction potential of copper is more positive than that of zinc.
- Copper ions would spontaneously oxidize aluminum.


- I only
- II only
- I and III
- I, II and III

17. Commercial aluminum is formed electrolytically from aluminum oxide (Al_2O_3), which is reduced at the cathode. Approximately how long must a current of 1000A be applied to form 50 g of aluminum?



- 1 second
- 1 1/2 minutes
- 10 minutes
- 17 hours

The following passage pertains to questions 18 - 22.

Thiamine pyrophosphate (TPP) is a thiamine (vitamin B1) derivative consisting of a pyrimidine ring which is connected to a thiazole ring, which is in turn connected to a pyrophosphate (diphosphate) functional group.




TPP performs an essential role in the catalytic activity of a number of enzymes. TPP is synthesized in the mitochondria for the activity of the pyruvate dehydrogenase complex (PDC). PDC is a complex of three enzymes that converts pyruvate into acetyl-CoA. This complex links glycolysis to the citric acid cycle. IN the mechanism of the pyruvate dehydrogenase enzyme, the anionic C2 carbon of TPP performs a nucleophilic attack on the C2 carbonyl of pyruvate. The resulting hemithioacetal undergoes decarboxylation.




The two carbon remnant of pyruvate, now attached to TPP, attacks the sulfur of the disulfide lipoic acid ring coenzyme attached to a lysine residue of the dihydrolipoyl transacetylase enzyme of the complex. In a ring-opening SN₂-like mechanism, the second sulfur is displaced as a sulfide. Release of the TPP cofactor and generates a thioacetate on lipoate. This is the rate-limiting step of the whole pyruvate dehydrogenase complex.

At this point, the lipoate-thioester functionality is translocated into the dihydrolipoyl transacetylase active site, where a transacylation reaction transfers the acetyl from the "swinging arm" of lipoyl to the thiol of coenzyme A. This produces acetyl-CoA, which is released from the enzyme complex and subsequently enters the citric acid cycle.



The dihydrolipoate, still bound to a lysine residue of the complex, then migrates to the dihydrolipoyl dehydrogenase active site where FAD converts dihydrolipoate back to its lipoate resting state, producing FADH₂. Then, a NAD⁺ cofactor converts FADH₂ back to its FAD resting state, producing NADH.



18. Which of the following describes the activity of NAD⁺ in the dihydrolipoyl dehydrogenase mechanism.

- I. oxidizing agent
- II. reducing agent
- III. Brønsted base

- A. I only
- B. II only
- C. I and III
- D. I, II and III

19. In the dihydrolipoyl transacetylase mechanism, the conversion of the disulfide into the dithiol form of lipoamide produces which change in the oxidation state of each sulfur atom?

- A. -1 to -2
- B. 0 to -1
- C. 0 to -2
- D. 0 to +1

20. Which is a net result of the mechanism of pyruvate dehydrogenase complex?
- A. Flavin is oxidized and nicotinamide is reduced.
 - B. Sulfur is oxidized and flavin is reduced.
 - C. Carbon is oxidized and nicotinamide is reduced.
 - D. Sulfur is oxidized and nicotinamide is reduced.
21. The standard reduction potential, E° , of lipoamide disulfide is -0.29V . The standard reduction potential of NAD^+ is -0.32V . Which of the following is necessarily true regarding the mitochondrial concentrations of these species during aerobic metabolism?
- A. $[\text{NAD}^+][\text{lipoamide dithiol}] > [\text{NADH}][\text{lipoamide disulfide}]$
 - B. $[\text{NAD}^+][\text{lipoamide disulfide}] > [\text{NADH}][\text{lipoamide dithiol}]$
 - C. $[\text{FAD}][\text{NADH}] > [\text{lipoamide dithiol}][\text{NAD}^+]$
 - D. $[\text{lipoamide dithiol}][\text{lipoamide disulfide}] > [\text{NAD}^+][\text{FAD}]$
22. The carbon atom of the carboxyl group of pyruvate will be oxidized, becoming CO_2 through the mechanism of the pyruvate dehydrogenase enzyme of the complex. What is the oxidizing agent?
- A. thiamine pyrophosphate
 - B. lipoamide dithiol
 - C. another carbon of pyruvate
 - D. coenzyme A

Redox & Electrochemistry

Answers and Explanations

1. A

Redox is an accounting system. The valence electrons in a molecular, atomic or ionic species are all assigned to an element. Electrons within a covalent bond are assigned to the more electronegative element. This accounting system provides a useful way to keep track of how electrons shift from their old neighborhoods in the reagents to their new neighborhoods in the products. An electron which before was in the vicinity of a lithium atom now finds itself in the vicinity of a fluorine atom. It has fallen down into a well. An oxidizing agent (oxidant, oxidizer) is a substance that has the ability to oxidize other substances — in other words to accept their electrons.

2. D

There are two tracks you can choose from in assigning oxidation numbers. You can use the rules for assigning oxidation numbers, or if you know or are given the structural formula, you can think of each bond as a tug of war and assign the electrons brought to the bonds to the more electronegative element.

In our molecule, each chlorine has brought seven valence electrons to share with oxygen in covalent bonding. Although they are both very electronegative elements, oxygen is more electronegative than chlorine (3.5 vs. 3.0), so the electrons chlorine brought to these bonds become oxygen's property in terms of redox accounting. The bond is only slightly polar, but we say those electrons are oxygen's. That's redox accounting. Oxygen has gained and chlorine has lost. Chlorine's oxidation number here is +7.

Alternatively, we could have used rules. One of our rules is that the oxidation number of oxygen is almost always -2 . Another rule is that the sum of oxidation numbers in an electrically neutral molecule is 0. We have seven oxygens. $7 \times -2 = -14$, so the two chlorines must be +7 each. Rules might be confusing though, because another rule is that the oxidation number of chlorine is almost always -1 . One of the exceptions is that when it is bonded to oxygen.

3. B


In the reaction, copper's oxidation number changes from +2 to 0. Iron's oxidation number changes from 0 to +2. Copper is reduced and iron is oxidized. Copper is the oxidizing agent and iron is the reducing agent.

4. A

In most reactions with O_2 , the oxygen atoms gain electron control over electrons brought by less electronegative elements to the reaction. In many reactions with H_2 hydrogen atoms lose electron control to more electronegative atoms. However, hydrogen's electronegativity is actually pretty respectable (2.1), so H_2 will not be readily oxidized by any metal.

5. D


It's very important to understand that hydrogen in its +1 oxidation state, such as in water or with H^+ ions, can perform as a decent oxidizing agent. It's easy to forget in the world of organic chemistry, where everything is oxidizing hydrogen, that hydrogen, itself, actually has a respectable electronegativity (2.1), especially compared to metals. If a very reactive, ie. low electronegativity, metal such as lithium is exposed to water, a vigorous reaction will result in which hydrogen oxidizes lithium.



The other metals of our answer choices are among the more electronegative transition metals. They are too electronegative to be oxidized by hydrogen.


6. B

Potassium permanganate is widely used as a strong oxidizing agent. The manganese atom has a high positive reduction potential. In a +7 oxidation state, its valence shell is all oxygen's property in $KMnO_4$. The manganese atom is reduced (+7 to +2) when potassium permanganate oxidizes a substrate, such as oxidation of acetaldehyde to acetic acid below.



7. B

A representative reaction of the type mentioned in the question stem would be sodium metal reacting with aluminum chloride to produce sodium chloride and aluminum. Aluminum oxidizes the sodium.



With electronegativity of 1.61, you would not normally expect to see aluminum oxidizing anything, but sodium's electronegativity is even lower, 0.93. In a tug-of-war with sodium over electrons, aluminum wins.

In other words, sodium is a great reducing agent. The best, most precise way to say that something is a powerful reducing agent is to say that its oxidized form, here sodium ion, has a large negative standard reduction potential.



This means that if you were removing electrons from a standard hydrogen electrode and trying to put them onto sodium ions, the electrons would have a long uphill climb to get there, 2.71 joules to climb up for every coulomb of electrons.

Think of the standard hydrogen electrode as the ground floor of a house. Some other places an electron might find itself are like the roof of the house. These are reducing agents like sodium, lithium or potassium. Reducing agents have negative standard reduction potential. Electrons have an uphill climb to get to these from the hydrogen electrode. Some other places are like the basement of the house, such as fluorine or oxygen. These are oxidizing agents. Oxidizing agents have positive standard reduction potential.


Aluminum itself has a substantial negative standard reduction potential.




Even though aluminum itself is a decent reducing agent, it will oxidize sodium because the trip for an electron from sodium to aluminum is downhill -2.71V to -1.66V.

8. B

Balancing by oxidation number method begins by determining the change in oxidation states of all of the individual species in the reaction and then balancing the increase with the decrease. The oxidation number decrease (minus 6 total), in chromium, is balanced with the increase, in chlorine (+6), when the coefficients are as follows.



The above is sufficient to answer the question, but let's look at the finished balanced reaction after inclusion of water and hydrogen ions.



Though balancing redox reactions is a classic Chem 101 problem, it's not a typical MCAT exercise. It's important to see and understand how this kind of thing works. It's important to understand that the oxidation number increase in a redox reaction must be matched by an oxidation number decrease.

9. C

Disproportionation is a redox reaction in which one compound of intermediate oxidation state converts to two compounds, one of higher and one of lower oxidation states. As an example, phosphorous acid disproportionates upon heating to give phosphoric acid and phosphine.



The reaction in the problem is the reverse of a disproportionation. A substance is formed, N_2 , in an intermediate oxidation state from precursors of lower and higher oxidation states. The reverse of disproportionation is called comproportionation.

10. B

Fluorine is an example of a substance with a large positive standard reduction potential. It is a powerful oxidizing agent.



This means that if you removed electrons from a standard hydrogen electrode and gave them to fluorine, the electrons would fall down into a deep well of binding energy, 2.87 joules for every coulomb of electrons.

11. A


Oxidation always occurs at the anode, whether the cell is electrolytic or galvanic. The reaction occurring in an electrolytic cell is non-spontaneous. An external voltage is being applied to make the reaction occur. What this means is that the electrons are being pushed uphill by the voltage, the overall cell potential being negative. Uphill for electrons is the journey from a positive potential to a negative potential. The external voltage is pulling the electrons out from the positive anode, where oxidation is occurring, and pushing them up onto the negative cathode, where reduction is occurring.

12. C

Oxidation always occurs at the anode, whether the cell is electrolytic or galvanic. The reaction occurring in an electrolytic cell is spontaneous. Electrons are falling downhill. The overall cell potential is positive. Downhill for electrons is the journey from a negative anode, where oxidation is occurring, to a positive cathode, where reduction is occurring.

13. D

Electrolysis of brine (concentrated NaCl) will oxidize Cl^- ions at the anode forming Cl_2 . The only other possible reductant would be the oxygen in water (hydrogen and sodium are already oxidized). Oxygen is generally more electron greedy than chlorine. In other words, it has a higher positive standard reduction potential than chlorine, so the voltage will pull the electrons from chloride preferentially over oxygen. Therefore, the anode half reaction is as follows.




At the cathode, the question of whether hydrogen or sodium will be reduced is easy. Sodium ions have a big negative standard reduction potential while hy-

drogen, of course, has a standard reduction potential of zero. It's the ground floor we compare everything to. Compared to hydrogen, sodium ions are a place it takes 2.71 joules of energy to move a coulomb of electrons to from a standard hydrogen electrode.



Compared to sodium, in other words, hydrogen is a very respectable oxidizing agent, so hydrogen is reduced at the cathode, not sodium.



In summary, electrolysis of brine produces chlorine gas at the anode and hydrogen gas at the cathode. As the reaction progresses, the solution forms concentrated sodium hydroxide.

14. C

The standard reduction potential represents the path that electrons would take from a standard hydrogen electrode to reduce a particular oxidant. The standard reduction potential of Zn^{2+} tells us that for hydrogen to reduce Zn^{2+} ions, electrons would need to travel uphill from hydrogen to -0.762V . To reduce Ag^+ the electrons would travel downhill, though, from hydrogen to $+0.800\text{V}$.

If you think about it, these two standard reduction potentials respectively have given us the two legs of the journey all the way from Zn (the reductant in our cell) to Ag^+ (the oxidant). Instead of going uphill to Zn^{2+} , we imagine going in the reverse, downhill instead from Zn to hydrogen. Instead of uphill to Zn^{2+} , downhill to hydrogen would be a change in the $+0.762\text{V}$ direction, and then from hydrogen to Ag^+ , which would be another $+0.800\text{V}$, so our cell potential is approximately 1.6V ($0.762\text{V} + 0.800\text{V}$). Don't let this be complicated. The electrons are falling from -0.762V to $+0.800\text{V}$.

The way you say this imaginary path as a formula is as follows.

$$E^{\circ}_{\text{cell}} = E^{\circ}_{\text{cathode}} - E^{\circ}_{\text{anode}}$$

15. D

One of the fundamental conceptual leaps we make into electrochemistry is to conceive of the free energy change in a chemical reaction as the journey of electrons through a potential difference. Instead of joules per mole we think in terms of joules per coulomb, or volts, as electrons transfer from the reductant to the oxidant. Underlying both conceptual frameworks is the fundamental reality that internal energy change in chemical reactions derive from the electrostatic potential energy differences as valence electrons find themselves within the new structural configurations of the product. Electrons were adjacent to a carbon atom and now they are in a covalent bond being pulled towards an oxygen nucleus, for example. For an oxidation-reduction reaction, we can convert a cell potential into a free energy change by using the Faraday unit for conversion of moles of electrons into Coulombs.

$$1 F = 96,500 \text{ C mol}^{-1}$$

$$\Delta G = -nFE$$

A very important equation in chemical thermodynamics describes how the free energy change of a reaction, ΔG , depends on both the standard free energy change, ΔG° , and the reaction quotient, Q .

$$\Delta G = \Delta G^\circ + RT \ln Q$$

In other words, whether a reaction is spontaneous in the forward or reverse direction depends not only on how the free energies of the product and reagent would compare, ΔG° , if they were both present in equal concentration. It also depends on what their concentrations actually are in the beaker, the reaction quotient, Q . A reaction with a negative standard free energy change, spontaneous forward if product and reagent are in equal concentration, may even run in the reverse direction if there is a high concentration of product and very little reagent.

We can convert this equation into a form which is convenient for electrochemistry. This new form of our thermodynamics equation is called the Nernst equation.

$$\Delta G = \Delta G^\circ + RT \ln Q$$

$$-nFE_{\text{cell}} = -nFE_{\text{cell}}^\circ + RT \ln Q$$

$$E_{\text{cell}} = E_{\text{cell}}^\circ - \frac{RT}{nF} \ln Q$$

$$E_{\text{cell}} = E_{\text{cell}}^\circ - \frac{0.0592V}{n} \ln Q$$

The actual cell potential depends not only on how the standard reduction potentials of the product and reagent sides of the reaction compare, E° , the cell potential if they were both present in equal concentration. The cell potential also depends on what their concentrations actually are in the electrochemical cell, which is given by the reaction quotient, Q .

As a redox reaction proceeds towards equilibrium, $Q \longrightarrow K$, the free energy change approaches zero, $\Delta G \longrightarrow 0$, and, equivalently, the cell potential approaches zero, $E_{\text{cell}} \longrightarrow 0$.

In summary, a dead battery is a chemical reaction that has attained the equilibrium state.

16. D

The evidence presented in the question stem is consistent with the relative standard reduction potentials of copper, zinc, and aluminum. On the table of standard reduction potentials below, we can see that aluminum metal is a stronger reducing agent than zinc metal which, in turn, is a stronger reducing agent than copper metal, and, equivalently, a copper ion is a stronger oxidizing agent than a zinc ion, which, in turn, is a stronger oxidizing agent than an aluminum ion. The standard reduction potential of a substance tells you the energy involved in reduction of the substance by the standard hydrogen electrode. The more negative the standard reduction potential, the more uphill the trip electrons must take in terms of potential energy. The more positive the standard reduction potential,

the further downhill the electrons are falling from hydrogen to carry out the reduction of the substance.

In the question stem, we were told that placing metallic zinc into a solution of 1M CuCl_2 will result in a layer of copper being deposited on the zinc. When metal ions are reduced, they 'plate out' as pure metal. This happens spontaneously because zinc is a stronger reducing agent than copper (choice I), ie. it has a more negative reduction potential than copper; copper has a more positive reduction potential than zinc (choice II).

Additionally, we were told that placing an aluminum rod in a 1M solution of ZnCl_2 will result in metallic zinc being deposited on the aluminum. This tells us that aluminum has a more negative reduction potential than zinc. If aluminum has a more negative reduction potential than zinc and zinc has a more negative reduction potential than copper, aluminum must through transitive principle have a more negative reduction potential than copper. It follows that copper ions would spontaneously oxidize aluminum.

E° (V)					
-2.93	K^+	$+ e^-$	\longrightarrow	K	
-2.71	Na^+	$+ e^-$	\longrightarrow	Na	
-1.66	Al^{3+}	$+ 3 e^-$	\longrightarrow	Al	
-0.76	Zn^{2+}	$+ 2 e^-$	\longrightarrow	Zn	
-0.44	Fe^{2+}	$+ 2 e^-$	\longrightarrow	Fe	
-0.25	Ni^{2+}	$+ 2 e^-$	\longrightarrow	Ni	
0	2H^+	$+ 2 e^-$	\longrightarrow	H_2	
+0.16	Cu^{2+}	$+ 2 e^-$	\longrightarrow	Cu	
+0.77	Fe^{3+}	$+ e^-$	\longrightarrow	Fe^{2+}	
+0.80	Ag^+	$+ e^-$	\longrightarrow	Ag	
+1.23	O_2	$+ 4 \text{H}^+$	$+ 4 e^-$	\longrightarrow	H_2O
+1.36	Cl_2	$+ 2 e^-$	\longrightarrow	Cl^-	
+1.51	MnO_4^-	$+ 8 \text{H}^+$	$+ 5 e^-$	\longrightarrow	$\text{Mn}^{2+} + 4 \text{H}_2\text{O}$
+2.08	O_3	$+ 2 \text{H}^+$	$+ 2 e^-$	\longrightarrow	$\text{O}_2 + 4 \text{H}_2\text{O}$
+2.87	F_2	$+ 2 e^-$	\longrightarrow	F^-	

17. C

It's unlikely to have a version of this type of problem on the MCAT involving extensive quantitative work, but you need to know how to get through a basic form. The key to relating the chemical stoichiometry of a redox reaction to DC current parameters is the conversion factor known as *the Faraday*.


$$1 F = 96,500 \text{ C mol}^{-1}$$

A mole of electrons is the same as 96,500 coulombs of electric charge.

We're making 50g of Al (s). The atomic weight of Al is 27 g mol^{-1} . Let's start rolling out conversion factors. This gets us to moles of aluminum.

$$50 \text{ g Al} \left(\frac{1 \text{ mol Al}}{27 \text{ g Al}} \right)$$

Aluminum in Al_2O_3 is in the +3 oxidation state. To plate out 1 mole of aluminum requires 3 moles of electrons.



This gets us to moles of electrons.

$$50 \text{ g Al} \left(\frac{1 \text{ mol Al}}{27 \text{ g Al}} \right) \left(\frac{3 \text{ mol } e^-}{1 \text{ mol Al}} \right)$$

Now the Faraday to get to electric charge.

$$50 \text{ g Al} \left(\frac{\text{mol Al}}{27 \text{ g Al}} \right) \left(\frac{3 \text{ mol } e^-}{\text{mol Al}} \right) \left(\frac{96,500 \text{ C}}{\text{mol } e^-} \right)$$

We were given that the current employed is 10,000A. If 1,000 coulombs flow per second, there is one second per 1,000 coulombs.

$$50 \text{ g Al} \left(\frac{\text{mol Al}}{27 \text{ g Al}} \right) \left(\frac{3 \text{ mol } e^-}{\text{mol Al}} \right) \left(\frac{96,500 \text{ C}}{\text{mol } e^-} \right) \left(\frac{1 \text{ s}}{1,000 \text{ C}} \right)$$

At the start, you don't need to see the whole solution to your foothold with a problem like this. Feel your way. We've worked through conversion factors all the way from grams of aluminum to the time required. Our answers are well spaced, so let's call $^{50}_{27} \sim 2$ and $96,500 \sim 100,000$.

$$50 \text{ g Al} \left(\frac{\text{mol Al}}{27 \text{ g Al}} \right) \left(\frac{3 \text{ mol } e^-}{\text{mol Al}} \right) \left(\frac{96,500 \text{ C}}{\text{mol } e^-} \right) \left(\frac{1 \text{ s}}{1,000 \text{ C}} \right)$$

$$\sim 600 \text{ s}$$

18. D

NAD^+ oxidizes FADH_2 . It is an oxidizing agent. Oxidation by NAD^+ always involves a two electron transfer in the form of a hydride (H^-).

Note that even though a hydrogen is transferred, it is not a proton transfer, but a hydride transfer. NAD^+ is not acting as a Brønsted base.

19. A

In lipoamide disulfide, each sulfur has one covalent bond to a carbon and one to a sulfur. Sulfur is slightly more electronegative than carbon (2.6 to 2.5), so the electron that carbon brought to that bond are assigned in redox accounting to sulfur. In lipoamide disulfide the oxidation state of sulfur is therefore -1 .

In the dithiol form of lipoamide, each sulfur has one covalent bond to a carbon and one to a hydrogen. Sulfur is more electronegative than both carbon and hydrogen, so each of the electrons those respective atoms brought to these bonds are now sulfur's property. In the dithiol form the oxidation state of sulfur is therefore -2 .

Not only in the context of lipoamide, but also regarding disulfide bridges in proteins and other important contexts such as the activity of glutathione, it's very important in biochemistry to recognize the disulfide form as the oxidized form and the dithiol as the reduced form.

20. C

As a result of the reactions of the pyruvate dehydrogenase complex, a pair of electrons that had been the property of carbon (originally within the nutrient molecule glucose) have become the property of nicotinamide. Between pyruvate and NADH , the pair of electrons were passed first to the two sulfurs of lipoamide, which then passed them to the flavin of FAD . FAD then passed them to NADH . Subsequent to completion of the mechanism, both lipoamide and FAD have returned to their initial oxidized state. In the net mechanism, carbon has been oxidized and NAD^+ reduced.

21. A

For of lipoamide disulfide to have a standard reduction potential of -0.29V means we would need to be pushing electrons uphill to NAD^+ at -0.32V , if the concentrations the reagents and products were all equal. In other words, the standard cell potential of a lipoamide/ NAD cell would be -0.03V .

$$E_{\text{cell}}^{\circ} = E_{\text{cathode}}^{\circ} - E_{\text{anode}}^{\circ}$$

$$-0.03\text{V} = (-0.32\text{V}) - (-0.29\text{V})$$

The reaction would not be spontaneous forward in that case. A negative standard cell potential corresponds to a positive standard free energy change.

$$\Delta G^{\circ} = -nFE^{\circ}$$

However, just as the actual free energy change not only depends on the standard free energy change but also the reaction quotient, so does the cell potential, which we can see in the Nernst equation.

$$\Delta G = \Delta G^{\circ} + RT \ln Q$$

$$-nFE_{\text{cell}} = -nFE_{\text{cell}}^{\circ} + RT \ln Q$$

$$E_{\text{cell}} = E_{\text{cell}}^{\circ} - \frac{RT}{nF} \ln Q$$

$$E_{\text{cell}} = E_{\text{cell}}^{\circ} - \frac{0.0592\text{V}}{n} \ln Q$$

For the dihydrolipoyl dehydrogenase mechanism to be spontaneous forward, the reaction quotient must be a fraction. The product of the concentrations of the reagents $[\text{NAD}^+][\text{lipoamide thiol}]$ must be greater than the product of the concentrations of the reaction products $[\text{NADH}][\text{lipoamide disulfide}]$.

22. C

Although a carbon of pyruvate is oxidized through the mechanism of the pyruvate dehydrogenase enzyme,

at this stage of the overall pyruvate dehydrogenase complex there has not yet been any net movement of electrons from the nutrient carbons to the electron shuttling coenzymes. The carboxyl carbon has lost an electron which the original carbonyl carbon of pyruvate has gained, so this is a type of disproportionation in which one carbon converts to a higher oxidation state and one to a lower oxidation state.

